



European Physical Society

HEP 2009

16-22 July 2009 Krakow, Poland



KLOE Measurement of the  $\sigma_{\pi\pi(\gamma)}$  cross section  
and the  $\pi^+\pi^-$  contribution to the muon anomaly




Federico Nguyen

INFN Roma TRE

for the KLOE Collaboration

Krakow - July, 17<sup>th</sup> 2009



# Outline

---

- Introduction: DAΦNE and KLOE
- Measurement of the  $\sigma_{\pi\pi(\gamma)}$  using ISR events with  $\gamma$  at small angle
- Determination of the  $\pi^+\pi^-$  contribution to  $a_\mu$
- Comparisons with recent  $e^+e^-$  experiments
- Outlook:  $\sigma_{\pi\pi(\gamma)}$  using ISR events with large angle  $\gamma$
- Conclusions




# DAΦNE and KLOE

$$L_{\text{peak}} = 1.3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$$

- ✓  $e^+ e^-$  collide at  $M_\phi$ :  $\sqrt{s} \sim 1.019 \text{ GeV}$
- ✓ angle btw the beams @ IP  $\sim 2 \times 12.5 \text{ mrad}$
- ✓ residual momentum in LAB  $\sim 13 \text{ MeV}$

2001-05:  $\sim 2.5 \text{ fb}^{-1}$  at  $M_\phi$

2006:  $\sim 250 \text{ pb}^{-1}$  at  $\sqrt{s}=1 \text{ GeV} + \sqrt{s}$  scan



Calorimeter, EmC:


Pb/Scint. Fiber, 4880 PMTs

98% of solid angle

$$\sigma_E / E = 0.057 / \sqrt{E} \text{ (GeV)}$$

$$\sigma_t = 57 \text{ ps} / \sqrt{E} \text{ (GeV)} \oplus 50 \text{ ps}$$

$$\sigma_{\perp} = 1.3 \text{ cm}$$



Drift Chamber, DC:

4 m  $\varnothing$   $\times$  3.3 m length

90% He, 10%  $i\text{-C}_4\text{H}_{10}$

12582 stereo sense wires

$$\sigma_p / p = 0.4\% \text{ for } \theta > 45^\circ$$

$$\sigma_{r\phi} = 0.150 \text{ mm}, \sigma_z = 2 \text{ mm}$$


$$\sigma(m_{\pi\pi}) \sim 1 \text{ MeV}$$

both detectors  
w/ trigger decision



# The cross section $\sigma_{e^+e^- \rightarrow \pi^+\pi^-}$ from ISR events

at a fixed  $\sqrt{s}$ , studying *Initial State Radiation* events,  $\sigma_{e^+e^- \rightarrow \pi^+\pi^-}(s)$  is extracted



$$\text{ISR only: } M_{\pi\pi}^2 \frac{d\sigma_{e^+e^- \rightarrow \pi^+\pi^-\gamma}}{dM_{\pi\pi}^2} = \sigma_{e^+e^- \rightarrow \pi^+\pi^-}(M_{\pi\pi}^2) \cdot H(M_{\pi\pi}^2, \theta_{\min})$$

→ EVA + PHOKHARA MC Generator

(S. Binner, J.H. Kühn, K. Melnikov, PLB459,1999)

(H.Czyż, A.Grzelińska, J.H Kühn, G.Rodrigo, EPJC27,2003)

**main advantage:**

no point-to-point errors on beam energy and luminosity

**main requirement:**

precise knowledge of ISR radiative corrections

1<sup>st</sup> KLOE publication (based on 140 pb<sup>-1</sup>)

A. Aloisio et al., PLB606(2005)12 → KLOE05

$$\frac{d\sigma_{\pi\pi\gamma}}{dM_{\pi\pi}^2} = \frac{N^{\text{obs}} - N^{\text{bkg}}}{\Delta M_{\pi\pi}^2} \cdot \frac{1}{\epsilon_{\text{sel}}} \cdot \frac{1}{L}$$





# Selection of $\pi\pi\gamma$ events at small angle

**PUBLISHED:**  
**PLB670(2009)285**

- a) 2 tracks with  $50^\circ < \theta_{\text{track}} < 130^\circ$
- b) small angle  $\gamma$  ( $\theta_{\pi\pi} < 15^\circ$ )

kinematics:  $\vec{p}_\gamma = \vec{p}_{\text{miss}} = -(\vec{p}_+ + \vec{p}_-)$

- ✓ high statistics for ISR ( $\sim \theta^{-2}$ )
- ✓ low relative FSR contribution
- ✓ suppressed  $\phi \rightarrow \pi^+\pi^-\pi^0$  wrt the signal



**statistics: 242pb<sup>-1</sup>**  
**3.4 Million Events**

# Selection of $\pi\pi\gamma$ events: suppress background



suppress  
 $e^+e^- \rightarrow e^+e^-\gamma$

$\pi/e$  separation performed with  
 particle ID based on the  
 calorimeter

remnant

$e^+e^- \rightarrow \mu^+\mu^-\gamma$  &  $\phi \rightarrow \pi^+\pi^-\pi^0$

cut and estimated as a function of  $M_{\pi\pi}^2$




$m_{\text{trk}}$ , defined under the  
 hypothesis of 2 equal  
 mass particles and 1  $\gamma$   
 in the final state



# Background estimates

Main backgrounds obtained from MC shapes fitted to data distribution in  $M_{\text{Trk}}$

$0.60 < M_{\pi\pi}^2 < 0.62 \text{ GeV}^2 \quad \chi^2/\text{ndf} = 158/180$




Data

$\Sigma$  MC

$\pi\pi\gamma$   $\mu\mu\gamma$

$\pi\pi\pi$   $ee\gamma$


$0.84 < M_{\pi\pi}^2 < 0.86 \text{ GeV}^2 \quad \chi^2/\text{ndf} = 179/258$




**Bkg errors are due to :**

- Uncertainty on  $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$  contribution

[EKHARA: PLB634\(2006\)493](#)




Bkg ( $\mu\mu\gamma$ ,  $\pi\pi\pi$  and  $ee\gamma$ ) fraction



- Error on normalization parameters obtained from the fit



# Data/MC corrections for the $\pi$ track

corrections obtained from  $\pi^+\pi^-\pi^0$  (large statistics, momentum limited) and  $\pi\pi\gamma$  data samples (after kin.  $\chi^2$ ) compatible within 0.3%  $\rightarrow$  syst. error




CAL

$$C_{3\pi} = 0.993 \pm 0.001$$



CAL

$$C_{\pi\pi\gamma} = 0.995 \pm 0.002$$






# Luminosity

KLOE measures  $L$  with Bhabha scattering

$55^\circ < \theta < 125^\circ$   
 acollinearity  $< 9^\circ$   
 $p \geq 400$  MeV

$$\int \mathcal{L} dt = \frac{N_{obs} - N_{bkg}}{\sigma_{eff}}$$



F. Ambrosino et al. (KLOE Coll.)  
**Eur.Phys.J.C47:589-596,2006**

generator used for  $\sigma_{eff}$   
**BABAYAGA (Pavia group)**

*C. M.C. Calame et al., NPB758 (2006) 22*

new version (**BABAYAGA@NLO**) gives  
 much better accuracy: 0.1%

Systematics on Luminosity	
Theory	0.1 %
Experiment	0.3 %
TOTAL 0.1 % th $\oplus$ 0.3% exp = 0.3%	



# Radiative corrections

- ISR-Process calculated at NLO-level

**PHOKHARA** generator (Czyż, Kühn et.al)

**Precision: 0.5%**

$$M_{\pi\pi}^2 \frac{d\sigma_{\pi\pi\gamma}}{dM_{\pi\pi}^2} = \sigma_{\pi\pi}(s) \times \mathbf{H}(s)$$



i) **Bare Cross Section**

divide by **Vacuum Polarisation**

→ from F. Jegerlehner




ii) **FSR - Corrections**

cross section  $\sigma_{\pi\pi}$  must be inclusive for FSR



FSR corrections taken into account in the efficiency evaluation and

in  $M_{\pi\pi}^2 \rightarrow M_{\gamma^*}^2$



# Systematic uncertainties


Systematic errors on  $a_\mu^{\pi\pi}$ :

Reconstruction Filter	negligible
Background	0.3%
$M_{\text{trk}}$ cuts	0.2%
$\pi/e$ ID and TCA	negligible
Tracking	0.3%
Hardware Trigger	0.1%
Acceptance ( $\theta_{\pi\pi}$ )	0.1%
Acceptance ( $\theta_\pi$ )	negligible
Unfolding	negligible
Software Trigger	0.1%
$\sqrt{s}$ dependence of H	0.2%
Luminosity( $0.1_{\text{th}} \oplus 0.3_{\text{exp}}$ )%	0.3%

experimental fractional error on  $a_\mu = 0.6\%$

FSR resummation	0.3%
Radiator H	0.5%
Vacuum polarization	0.1%


theoretical fractional error on  $a_\mu = 0.6\%$



$\sigma_{\pi\pi}$ , undressed from VP, inclusive for FSR as function of  $(M_{\gamma^*})^2$



# Recent comparisons on $a_{\mu}^{\pi\pi}$




$$a_{\mu}^{\pi\pi} = \frac{1}{4\pi^3} \int_{0.35\text{GeV}^2}^{0.95\text{GeV}^2} ds \sigma(e^+e^- \rightarrow \pi^+\pi^-) K(s)$$

average value in  $10^{-10}$  units:


$359.2 \pm 2.1$  with  $\chi^2/\text{dof} = 1.24/2$   
 confidence level of 54%

even at the largest difference, minor impact on  $a_{\mu}^{\pi\pi}$

*$e^+e^-$  experiments are consistent*



# Present situation on $a_\mu$





KLOE strengthens the discrepancy  $\sim 3.4 \sigma$  between the SM prediction and the BNL measurements



# NEW: selection of ISR $\gamma$ at large angle


- ✓ independent complementary analysis
- ✓ threshold region  $(2m_{\pi})^2$  accessible
- ✓  $\gamma_{\text{ISR}}$  photon detected  
(4-momentum constraints)
- ✓ background from  $\phi$  decays,  $\phi \rightarrow \pi^+\pi^-\pi^0$   
&  $\phi \rightarrow f_0(980)\gamma \rightarrow \pi\pi\gamma$  suppressed using  
data taken at  $\sqrt{s} = 1$  GeV, off the  $\phi$  peak

detection of 2 tracks and  
at least 1  $\gamma$  ( $E > 50$  MeV)





Federico Nguyen  
17-07-2009

# Control of backgrounds: $A_{FB}$



$$C_{\pi\pi} = -1$$


$A(\text{FSR}), A(\text{S}\gamma), A(\rho\pi)$   
are C-even



$A_{FB}$  more sensitive than  $d\sigma/dM$

MC based on  
PHOKHARA: Phys.Lett.B611(2005)116  
FASTERD: arXiv:0901.4440 [hep-ph]

*accurate control of same final  
state interfering backgrounds*



# Conclusions

---




- ✓ we presented a new measurement of  $\sigma_{\pi\pi(\gamma)}$  and of the  $\pi^+\pi^-$  contribution to  $a_\mu^{\pi\pi}$  in the range  $[0.35, 0.95] \text{ GeV}^2$  with 0.9% accuracy [PLB670 (2009) 285]
- ✓ this result is in good agreement with the CMD-2 and SND recent results, and it strengthens the difference between BNL measurement and SM prediction
- ✓ an independent analysis with  $\gamma$  detected at large angle is very close to be finalized (selection cuts established and main corrections evaluated), preliminary data-MC comparison shows excellent agreement
- ✓ we plan to determine the  $\pi^+\pi^-$  contribution to  $a_\mu^{\pi\pi}$ , from ratio of  $\pi\pi\gamma$  to  $\mu\mu\gamma$  events, that allows an independent check of the radiator function and cancellation of some systematic effects





# Comparisons on $F_\pi$


$$\sigma_{\pi\pi} = \frac{\pi\alpha^2\beta_\pi^3}{3s} |F_\pi|^2$$



*good agreement below and on the  $\rho$  peak among different  $e^+e^-$  experiments*




# Trigger corrections



average value =  $0.9987 \pm 0.0002$

the main source (hardware veto of cosmic rays) of inefficiency in the 2005 result has been removed





trigger efficiency:  
fractional error given by  
relative difference of 2  
independent methods  
from data  $\rightarrow$  0.1%







# Geometrical acceptance for the $\gamma$

we study the impact of varying the  $15^\circ$  cut on  $\theta_{\pi\pi}$  in slices of  $M^2_{\pi\pi}$

the data/MC spectrum variation is linear as a function of the cut, so the excursion at  $\pm 1$  degree is taken as systematic error



# K LOng Experiment: resolutions



Federico Nguyen  
17-07-2009

# Electron/pion identification

$\pi/e$  separation is performed using a particle ID function based on time and quantity and shape of the energy released in the calorimeter

