

a “mass preconditioning” for lattice Dirac operators

G.M. de Divitiis^{a,b}, R. Petronzio^{a,b}, N. Tantalo^{a,c}

^a INFN sez. “Tor Vergata”

^b Rome University “Tor Vergata”

^c Centro Ricerche e Studi “E. Fermi”

07-05-2010

- a numerical problem: heavy quark propagators decay too fast!
 - the free theory case
 - the interacting case

- a numerical trick: heavy quark propagators decay as slow as light quark propagators
 - the free theory case
 - the interacting case
 - different preconditioning

- the same numerical trick: speeding up numerical inversions for light quark

a numerical problem: heavy quark propagators decay too fast!

first principle approaches to flavour physics are of fundamental importance in the search for physics beyond the Standard Model

in order to calculate heavy flavour observables on the lattice we need to solve the linear system

$$(D + M) \psi = \eta^{y_0}$$


on the one hand, the numerical inversion is quite fast for heavy quarks

on the other hand, at large times the solution is poorly accurate because $|\psi(x_0; \vec{x})|$ may become much smaller than r for $x_0 \gg y_0$

a solver:

- $n \rightarrow n + 1$
- apply $(D + M)$ a few times ...
- a bit of linear algebra ...
- check if $|(D + M) \psi^n - \eta^{y_0}| < r$



a numerical problem: free theory




- this is a “heavy” pseudoscalar-pseudoscalar correlator in free theory
- we do expect that if we choose a residue that is too big something should go wrong at large time distances from the source ...
- in this particular case we can compare the numerical inversion performed with a “big” residue $r = 10^{-6}$ with the one performed with a “small” residue $r = 10^{-11}$

a numerical problem: free theory

$M_{PP}(t)$




a numerical problem: free theory




- by changing a bit the quark mass the effect may become “particularly” evident
- in this work we have analyzed situations in which the problem can be easily identified and such that “exact” results can be obtained by working with double precision architecture
- we have been working with “big” residues $r \sim 10^{-6}$ also because this is the best one can do with single precision architectures (GPU are much faster in single precision)

a numerical problem: interacting theory



- here we see the same effect in the interacting theory...
- simulation details: $\beta = 5.3$, $k_{sea} = 0.13625$, $am_{sea} \simeq 0.07$, $am_t \simeq 0.35$
- we have been working with "big" residues $r \sim 10^{-6}$ also because this is the best one can do with single precision architectures (GPU are much faster in single precision)

a numerical trick: free theory




we choose:

$$\psi(\vec{x}, x_0) \longrightarrow \cosh [m_0(x_0 - T/2)] \chi(\vec{x}, x_0)$$

and, in this particular case, $m_0 = 0.4$. We calculate numerically $\chi(\vec{x}, x_0)$.

a numerical trick: free theory




we choose:

$$\psi(\vec{x}, x_0) \longrightarrow \cosh [m_0(x_0 - T/2)] \chi(\vec{x}, x_0)$$

and, in this particular case, $m_0 = 0.4$. We calculate numerically $\chi(\vec{x}, x_0)$, and *offline* we get $\psi(\vec{x}, x_0)$

a numerical trick: free theory




we choose:

$$\psi(\vec{x}, x_0) \longrightarrow \cosh [m_0(x_0 - T/2)] \chi(\vec{x}, x_0)$$

and, in this particular case, $m_0 = 0.4$. We calculate numerically $\chi(\vec{x}, x_0)$, and *offline* we get $\psi(\vec{x}, x_0)$

a numerical trick: interacting theory



also in the interacting case we choose:


$$\psi(\vec{x}, x_0) \longrightarrow \cosh [m_0(x_0 - T/2)] \chi(\vec{x}, x_0)$$

and $m_0 = 0.4$. We calculate numerically $\chi(\vec{x}, x_0)$, and *offline* we get $\psi(\vec{x}, x_0)$


a numerical trick: Schrödinger Functional setup

up to now we have been discussing the case of periodic boundary in the time direction

our preconditioning may be particularly relevant in the case of fixed boundary conditions in the time direction:



a numerical trick: SF free theory




we choose:

$$\psi(\vec{x}, x_0) \longrightarrow \exp(m_0 x_0) \chi(\vec{x}, x_0)$$

We calculate numerically $\chi(\vec{x}, x_0)$, and *offline* we get $\psi(\vec{x}, x_0)$

a numerical trick: SF free theory




we choose:

$$\psi(\vec{x}, x_0) \longrightarrow \exp(m_0 x_0) \chi(\vec{x}, x_0)$$

We calculate numerically $\chi(\vec{x}, x_0)$, and *offline* we get $\psi(\vec{x}, x_0)$

a numerical trick: SF interacting theory



- here we see the same effect in the interacting theory...
- simulation details: $\beta = 6.638$, $L \sim 0.5$ fm
- this is a "small volume" simulation needed to study B -physics on the lattice by using the so-called Step Scaling Method
- with this preconditioning it will be possible to separate "genuine" finite volume effects from "excited states contaminations" by making small volume simulations with $T = 4L$


a numerical trick: different preconditioning

- the preconditioning that we have been discussing up to now
 - it is particularly simple to implement
 - solves the large time numerical precision issue for heavy quark propagators,
 - can be “removed” after having computed the correlation functions
- by relaxing the last property, one can as easily as before explore several other possibilities
 - extend the same trick to the other directions
 - give to the matrix α a “structure” in Dirac space
 - ...

in the following we shall briefly discuss the following preconditioning:

$$\psi(x_0, x_1, x_2, x_3) \longrightarrow \alpha(x_0) \alpha(x_1) \alpha(x_2) \alpha(x_3) \chi(x_0, x_1, x_2, x_3)$$


a numerical trick: different preconditioning



we choose ($m_0 = 0.4$):

$$\psi(x_0, x_1, x_2, x_3) \longrightarrow \left(\prod_{i=0}^3 \cosh [m_0(x_i - L_i/2)] \right) \chi(x_0, x_1, x_2, x_3)$$

a numerical trick: different preconditioning



we get the same answer as before but by performing many more iterations for the solver to converge:

- simulation details: $\beta = 5.3$, $k_{sea} = 0.13625$, $am_{sea} \simeq 0.07$, $am_t \simeq 0.35$
- DD-fgcr inveter $r = 10^{-11}$ unpreconditioned: 7 iterations
- DD-fgcr inveter $r = 10^{-6}$ time preconditioned: 6 iterations
- DD-fgcr inveter $r = 10^{-6}$ all- d preconditioned: 23 iterations

the same numerical trick: speeding up numerical inversions for light quarks

we make light quark propagators decay faster by choosing:

$$\psi(x_0, x_1, x_2, x_3) \longrightarrow \left(\prod_{i=0}^3 \frac{1}{\cosh [m_0(x_i - L_i/2)]} \right) \chi(x_0, x_1, x_2, x_3)$$

| | β | $L^3 \times T$ | k_{sea} | r | m_0 | iterations |
|-----------|---------|------------------|-----------|------------|-------|------------|
| <i>D5</i> | 5.3 | $24^3 \times 48$ | 0.13625 | 10^{-11} | 0.0 | 175 |
| <i>D5</i> | 5.3 | $24^3 \times 48$ | 0.13625 | 10^{-11} | 0.4 | 141 |
| <i>E3</i> | 5.3 | $32^3 \times 64$ | 0.13605 | 10^{-10} | 0.0 | 100 |
| <i>E3</i> | 5.3 | $32^3 \times 64$ | 0.13605 | 10^{-10} | 0.2 | 79 |
| <i>E3</i> | 5.3 | $32^3 \times 64$ | 0.13605 | 10^{-10} | 0.4 | 70 |
| <i>E4</i> | 5.3 | $32^3 \times 64$ | 0.13610 | 10^{-10} | 0.0 | 115 |
| <i>E4</i> | 5.3 | $32^3 \times 64$ | 0.13610 | 10^{-10} | 0.2 | 91 |
| <i>E4</i> | 5.3 | $32^3 \times 64$ | 0.13610 | 10^{-10} | 0.4 | 81 |
| <i>E5</i> | 5.3 | $32^3 \times 64$ | 0.13625 | 10^{-10} | 0.0 | 194 |
| <i>E5</i> | 5.3 | $32^3 \times 64$ | 0.13625 | 10^{-10} | 0.2 | 153 |
| <i>E5</i> | 5.3 | $32^3 \times 64$ | 0.13625 | 10^{-10} | 0.4 | 141 |

- we have considered a “family” of preconditioning that are easy to implement
- that can be used to perform “flavored” quark inversions on single precision architectures (e.g. GPUs, Cell, etc.) with the same numerical accuracy one would get on n -precision architectures
- on double precision machines, our preconditioning can be used to speed up the calculation of light quark propagators
- we have demonstrated that one can easily gain up to 30% in computational time without compromising the numerical accuracy
- we are working to use such kind of preconditioning in the HMC generation of gauge field configurations
- we are exploring several other possibilities with respect to the ones discussed in this talk and, in particular, giving a Dirac “structure” to the preconditioning operator